

**Motor-Vehicle Occupant Injury:
Strategies for Increasing Use
of Child Safety Seats, Increasing Use
of Safety Belts, and
Reducing Alcohol-Impaired Driving**

**A Report on Recommendations of the Task
Force on Community Preventive Services**

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
Centers for Disease Control and Prevention (CDC)
Atlanta, GA 30333



The *MMWR* series of publications is published by the Epidemiology Program Office, Centers for Disease Control and Prevention (CDC), U.S. Department of Health and Human Services, Atlanta, GA 30333.

SUGGESTED CITATION

Centers for Disease Control and Prevention. Motor-vehicle occupant injury: strategies for increasing use of child safety seats, increasing use of safety belts, and reducing alcohol-impaired driving. A report on recommendations of the Task Force on Community Preventive Services. *MMWR* 2001;50(No. RR-7):[inclusive page numbers].

Centers for Disease Control and Prevention Jeffrey P. Koplan, M.D., M.P.H.
Director

The production of this report as an *MMWR* serial publication was coordinated in
Epidemiology Program Office Stephen B. Thacker, M.D., M.Sc.
Director

Office of Scientific and Health Communications John W. Ward, M.D.
Director
Editor, MMWR Series

Recommendations and Reports Suzanne M. Hewitt, M.P.A.
Managing Editor

Amanda Crowell
Project Editor

Lynda G. Cupell
Visual Information Specialist

Michele D. Renshaw
Erica R. Shaver
Information Technology Specialists

Contents

Background	1
Introduction	2
Methods	2
Results	5
Use of the Recommendations in States and Communities	6
Additional Information Regarding the <i>Community Guide</i>	11
References	12

Task Force on Community Preventive Services*

CHAIR

Caswell A. Evans, Jr., D.D.S, M.P.H.
National Institute for Dental and
Craniofacial Research
National Institutes of Health
Bethesda, Maryland

VICE-CHAIR

Jonathan E. Fielding, M.D., M.P.H., M.B.A.
Los Angeles Department of Health Services
Los Angeles, California

MEMBERS

Ross C. Brownson, Ph.D.
St. Louis University School of Public Health
St. Louis, Missouri

Mary Jane England, M.D.
Washington Business Group on Health
Washington, DC

Mindy Thompson Fullilove, M.D.
New York State Psychiatric Institute and
Columbia University
New York, New York

Fernando A. Guerra, M.D., M.P.H.
San Antonio Metropolitan Health District
San Antonio, Texas

Alan R. Hinman, M.D., M.P.H.
Task Force for Child Survival and
Development
Atlanta, Georgia

George J. Isham, M.D.
Health Partners
Minneapolis, Minnesota

Garland H. Land, M.P.H.
Center for Health Information Management
and Epidemiology
Missouri Department of Health
Jefferson City, Missouri

Charles S. Mahan, M.D.
College of Public Health
University of South Florida
Tampa, Florida

Patricia Dolan Mullen, Dr.P.H.
University of Texas-Houston
School of Public Health
Houston, Texas

Patricia A. Nolan, M.D., M.P.H.
Rhode Island Department of Health
Providence, Rhode Island

Susan C. Scrimshaw, Ph.D.
School of Public Health
University of Illinois
Chicago, Illinois

Steven M. Teutsch, M.D., M.P.H.
Merck & Company, Inc.
West Point, Pennsylvania

Robert S. Thompson, M.D.
Department of Preventive Care
Group Health Cooperative of Puget Sound
Seattle, Washington

*Patricia A. Buffler, Ph.D., M.P.H., University of California, Berkeley, and David W. Fleming, M.D., CDC, Atlanta, Georgia, served on the Task Force while most of the recommendations were being developed.

The following CDC staff members prepared this report:

Stephanie Zaza, M.D., M.P.H.

Kate W. Harris

Patryce V. Young-Curtis, M.S.H.C.A.

Erin P. Finley

*Division of Prevention Research and Analytic Methods
Epidemiology Program Office*

Ruth A. Shults, Ph.D., M.P.H.

David A. Sleet, Ph.D., M.A.

Randy W. Elder, M.Ed.

Tho Bella Dinh-Zarr, M.P.H.

*Division of Unintentional Injury Prevention
National Center for Injury Prevention and Control*

Daniel M. Sosin, M.D., M.P.H.

Office of the Director

National Center for Injury Prevention and Control

in collaboration with

James L. Nichols, Ph.D.

Office of Research and Traffic Records

Traffic Safety Programs

National Highway Traffic Safety Administration

Robert S. Thompson, M.D.

Task Force on Community Preventive Services

and

Group Health Cooperative of Puget Sound

Motor-Vehicle Occupant Injury: Strategies for Increasing Use of Child Safety Seats, Increasing Use of Safety Belts, and Reducing Alcohol-Impaired Driving

A Report on Recommendations of the Task Force on Community Preventive Services

Summary

The Task Force on Community Preventive Services has conducted systematic reviews of interventions designed to increase use of child safety seats, increase use of safety belts, and reduce alcohol-impaired driving. The Task Force strongly recommends the following interventions: laws requiring use of child safety seats, distribution and education programs for child safety seats, laws requiring use of safety belts, both primary and enhanced enforcement of safety belt use laws, laws that lower the legal blood alcohol concentration (BAC) limit for adult drivers to 0.08%, laws that maintain the minimum legal drinking age at 21 years, and use of sobriety checkpoints. The Task Force recommends communitywide information and enforcement campaigns for use of child safety seats, incentive and education programs for use of child safety seats, and a lower legal BAC for young drivers (in the United States, those under the minimum legal drinking age). This report provides additional information regarding these recommendations, briefly describes how the reviews were conducted, and provides information to help apply the interventions locally.

BACKGROUND

Motor-vehicle-related injuries kill more children and young adults (i.e., those aged 1–24 years) than any other single cause in the United States (1,2) and are the leading cause of death from unintentional injury for persons of all ages (3,4). Approximately 41,000 persons in the United States die in motor-vehicle crashes each year (5). Moreover, crash injuries result in approximately 500,000 hospitalizations and 4 million emergency department visits annually (6).

Viewed from an economic perspective, crash injuries and deaths are a burden to society. Motor-vehicle-related deaths and injuries cost the United States approximately \$150 billion annually (7,8), including \$52.1 billion in property damage, \$42.4 billion in lost productivity, and \$17 billion in medical expenses (7). Alcohol-related crashes contribute substantially to these costs, with a direct economic impact of approximately \$45 billion in 1994 alone (7).

Reducing motor-vehicle injury remains a formidable public health challenge, despite sharp declines in motor-vehicle-related death rates since 1925 (9). Use of child safety seats and safety belts and deterrence of alcohol-impaired driving are among the most important preventive measures to further reduce motor-vehicle occupant injuries

and deaths (10,11). This report provides recommendations on interventions to increase use of child safety seats, increase use of safety belts, and reduce alcohol-impaired driving.

INTRODUCTION

This *MMWR* report is the third to be completed for the *Guide to Community Preventive Services* (the *Community Guide*), a resource that will include multiple chapters, each focusing on a preventive health topic. The first two reports were on vaccine-preventable diseases and tobacco use prevention and reduction (12–17). This report provides an overview of the process used by the Task Force on Community Preventive Services (the Task Force) to select and review evidence and summarizes the recommendations of the Task Force regarding interventions to reduce motor-vehicle occupant injury. A full report of the recommendations, supporting evidence (i.e., discussions of applicability, additional benefits, potential harms, existing barriers to implementation, and economic evaluations), and remaining research questions will be published in the *American Journal of Preventive Medicine* later this year.

The independent, nonfederal Task Force is developing the *Community Guide* with the support of the U.S. Department of Health and Human Services (DHHS) in collaboration with public and private partners. CDC provides staff support to the Task Force for development of the *Community Guide*. The recommendations in this report, although developed independently by the Task Force, are consistent with current CDC recommendations.

METHODS

The *Community Guide's* methods for conducting systematic reviews and linking evidence to recommendations have been described elsewhere (18). In brief, for each *Community Guide* topic, a multidisciplinary team conducts a review by

- developing an approach to organizing, grouping, and selecting the interventions;
- systematically searching for and retrieving evidence;
- assessing the quality of and summarizing the strength of the body of evidence of effectiveness;
- summarizing information regarding other evidence; and
- identifying and summarizing research gaps.

For motor-vehicle occupant injury, the development team focused on interventions to promote use of child safety seats, promote use of safety belts, and deter alcohol-impaired driving. These areas were chosen because a) use of child safety seats and use of safety belts are below national goals (19); b) 38% of traffic deaths still involve alcohol (5); and c) nonuse of child safety seats, nonuse of safety belts, and alcohol-impaired driving are among the most important contributors to motor-vehicle occupant injuries, and reducing these three risk behaviors could dramatically reduce these injuries. This report includes the goals of the National Highway Traffic Safety Administration (NHTSA) and the *Healthy People 2010* initiative (19) in these areas (Table 1).

TABLE 1. Selected National Highway Traffic Safety Administration* (NHTSA) goals and *Healthy People 2010* objectives related to motor-vehicle occupant injury

NHTSA goal	<i>Healthy People 2010</i> objective [†]
General	
Reduce the number of fatal and nonfatal injuries by 20% by the year 2008 (from 42,065 fatal injuries and 3,511,000 nonfatal injuries in 1996). [§]	Reduce deaths caused by motor-vehicle crashes from 15.0/100,000 persons (1998 preliminary data age-adjusted to the year 2000 standard population) to 9.0/100,000. (Objective 15-15a)
	Reduce deaths from 2/100 million vehicle miles traveled (in 1997) to 1/100 million vehicle miles traveled. (Objective 15-15b)
	Reduce nonfatal injuries caused by motor-vehicle crashes from 1,270/100,000 persons (in 1997) to 1,000/100,000 (21% improvement). (Objective 15-17).
Child Safety Seats	
Reduce child (aged 0–4 years) occupant fatalities by 25% by the year 2005 (from 653 fatalities in 1996). [¶]	Increase use of child restraint devices for passengers aged 0–4 years from 92% (1998 preliminary data age-adjusted to the year 2000 standard population) to 100%. (Objective 15-20)
Safety Belts	
Increase national seat belt use to 90% by the year 2005 (from 68% in 1996). ^{**}	Increase use of safety belts from 69% (in 1998) to 92% (33% improvement). (Objective 15-19)
Alcohol-Impaired Driving	
Reduce alcohol-related fatalities to \leq 11,000 annually by the year 2005. ^{††} (Achieving this goal will reduce deaths caused by drinking and driving by approximately 5,000 each year.)	Reduce deaths caused by alcohol-related motor-vehicle crashes from 6.1/100,000 persons (1997 baseline) to 4/100,000. (Objective 26-1a)
	Reduce injuries caused by alcohol-related motor-vehicle crashes from 122/100,000 persons (1997 baseline) to 65/100,000. (Objective 26-1b)
	Reduce the proportion of adolescents who report that during the preceding 30 days they rode with a driver who had been drinking alcohol from 37% (in 1997) to 30%. (Objective 26-6)
	Extend administrative license revocation laws or programs of equal effectiveness for persons who drive under the influence of intoxicants from 41 states (in 1998) to all states and Washington, D.C. (Objective 26-24)
	Extend legal requirement for maximum blood alcohol concentration levels of 0.08% for motor-vehicle drivers aged \geq 21 years from 16 states (in 1998) to all states and Washington, D.C. (Objective 26-25)

* U.S. Department of Transportation.

[†] US Department of Health and Human Services. *Healthy People 2010*, vols I and II. 2nd ed. Washington, DC: US Government Printing Office, November 2000.[§] 1998 NHTSA Strategic Plan, available on the Internet at <<http://www.nhtsa.dot.gov/nhtsa/whatis/planning/StratPlan.1998/>>.[¶] NHTSA. Presidential Initiative for Increasing Seat Belt Use Nationwide: Recommendations from the Secretary of Transportation. Washington, DC: US Department of Transportation, NHTSA, 1997; publication no. DOT HS 808 576.^{**} Presidential Initiative to Increase Seat Belt Use Nationwide, available at <<http://www.nhtsa.dot.gov/people/injury/airbags/buckleplan/presbelt2/#2>>.^{††} NHTSA Alcohol-Impaired Driving Goals: NHTSA Impaired Driving Division, available at <http://www.nhtsa.dot.gov/people/outreach/safesobr/ydydyl/imp_drive.html>.

The consultation team* generated a comprehensive list of strategies and created a priority list of interventions for review based on a process of polling consultants and other specialists in the field regarding their perception of the importance and practicality of various interventions.

Interventions reviewed were either single-component (i.e., using only one activity to achieve desired outcomes) or multicomponent (i.e., using more than one related activity). Studies were grouped on the basis of the similarity of the interventions being evaluated. Some studies provided evidence for more than one intervention. In these cases, the studies were reviewed for each applicable intervention. Interventions and outcome measures were classified according to definitions developed as part of the review process. The nomenclature used here might differ from that used in the original studies.

To be included in the reviews of effectiveness, studies had to a) be primary investigations of interventions selected for evaluation rather than, for example, guidelines or reviews; b) be published in English during 1966–June 2000; c) be conducted in established market economies;† and d) compare outcomes among groups of persons exposed to the intervention with outcomes among groups of persons not exposed or less exposed to the intervention (whether the comparison was concurrent between groups or before-after within groups).

For each intervention reviewed, the team developed an analytic framework indicating possible causal links between the intervention under study and predefined outcomes of interest. To make recommendations, the Task Force required that studies show increases in use of child safety seats or safety belts, decreases in alcohol-impaired driving, or decreases in motor-vehicle crashes or crash-related injuries.

*Members of the consultation team were Julie C. Bolen, Ph.D., M.P.H., National Center for Chronic Disease Prevention and Health Promotion, CDC, Atlanta, Georgia; Robert D. Brewer, M.D., M.S.P.H., Nebraska Department of Health, Lincoln, Nebraska; Stephanie D. Bryn, M.P.H., Health Resources Services Administration, Rockville, Maryland; Forrest M. Council, Ph.D., University of North Carolina, Chapel Hill, North Carolina; Robert W. Denniston, M.A., Substance Abuse and Mental Health Services Administration, Rockville, Maryland; Andrea C. Gielen, Sc.D., Sc.M., Johns Hopkins University, Baltimore, Maryland; Sue Gorcowski, M.A., National Highway Traffic Safety Administration, Washington, D.C.; Charles A. Hurley, National Safety Council, Washington, D.C.; Bruce H. Jones, M.D., M.P.H., National Center for Injury Prevention and Control, CDC, Atlanta, Georgia; Trudy A. Karlson, Ph.D., University of Wisconsin, Madison, Wisconsin; Mark R. Kinde, M.P.H., Minnesota Department of Health, Minneapolis, Minnesota; David W. Lawrence, M.P.H., San Diego State University, San Diego, California; Sue E. Martin, Ph.D., National Institute for Alcohol Abuse and Alcoholism, Rockville, Maryland; Jim A. McKnight, Ph.D., National Public Service Research Institute, Landover, Maryland; Angela D. Mickalide, Ph.D., National SAFE KIDS Campaign, Washington, D.C.; James L. Nichols, Ph.D., National Highway Traffic Safety Administration, Washington, D.C.; Lloyd F. Novick, M.D., M.P.H., Onondaga County Department of Health, Syracuse, New York; Fred P. Rivara, M.D., M.P.H., University of Washington, Seattle, Washington; Carol W. Runyan, Ph.D., M.P.H., University of North Carolina, Chapel Hill, North Carolina; Richard J. Smith, M.S., Indian Health Service, Rockville, Maryland; Patricia F. Waller, Ph.D., University of Michigan, Ann Arbor, Michigan; Allan F. Williams, Ph.D., Insurance Institute for Highway Safety, Arlington, Virginia.

† Established market economies as defined by the World Bank are Andorra, Australia, Austria, Belgium, Bermuda, Canada, Channel Islands, Denmark, Faeroe Islands, Finland, France, Former Federal Republic of Germany, Germany, Gibraltar, Greece, Greenland, Holy See, Iceland, Ireland, Isle of Man, Italy, Japan, Liechtenstein, Luxembourg, Monaco, The Netherlands, New Zealand, Norway, Portugal, San Marino, Spain, St. Pierre and Miquelon, Sweden, Switzerland, the United Kingdom, and the United States.

Improvements in behavioral outcomes (i.e., use of child safety seats, use of safety belts, and decreases in alcohol-impaired driving) are acceptable because

- child safety seats are 55%–70% effective in preventing deaths (20);
- safety belts are 45%–60% effective in reducing deaths and 50%–65% effective in reducing moderate-to-critical injuries (21); and
- the risk for fatal crash involvement increases as blood alcohol levels increase (22).

Each study that met the inclusion criteria was evaluated using a standardized abstraction form and assessed for suitability of the study design and threats to validity. On the basis of the number of threats to validity, studies were characterized as having good, fair, or limited execution (18,23). Results on each outcome of interest were obtained from each study that met the minimum quality criteria. For studies that reported multiple measures of a given outcome, the “best” measure with respect to validity and stability was chosen according to consistently applied rules. Measures that were adjusted for the effects of potential confounders were used in preference to crude effect measures. For studies in which such adjusted results were not provided, net effects were derived when possible by calculating the difference between the changes observed in the intervention and comparison groups. A median was calculated as a summary effect measure for each outcome of interest. For bodies of evidence consisting of seven or more studies, an interquartile range is presented as an index of variability; otherwise, a simple range is reported.

The strength of the body of evidence of effectiveness was characterized as strong, sufficient, or insufficient on the basis of the number of available studies, the suitability of study designs for evaluating effectiveness, the quality of execution of the studies, the consistency of the results, and the effect size (18).

The *Community Guide* uses systematic reviews to evaluate the evidence of intervention effectiveness, and the Task Force makes recommendations based on the findings of these reviews (18). The strength of each recommendation is based on the strength of the evidence of effectiveness (e.g., an intervention is “strongly recommended” when there is strong evidence of effectiveness or “recommended” when there is sufficient evidence) (18). Other types of evidence can also affect a recommendation. For example, evidence of harms resulting from an intervention might lead to a recommendation that the intervention not be used if adverse effects outweigh improved outcomes. In general, the Task Force does not use economic information to modify recommendations.

A finding of insufficient evidence of effectiveness should not be seen as evidence of ineffectiveness. Such a finding is important for identifying areas of uncertainty and continuing research needs. In contrast, adequate evidence of ineffectiveness leads to a recommendation that the intervention not be used.

RESULTS

Searches of six computerized databases (i.e., Medline, Embase, El Compindex, Sociological Abstracts, Psychlit, and Transportation Research Information Services [TRIS]*) yielded a list of 10,948 titles, from which 3,653 articles were retrieved as possibly

*These databases can be accessed through the Dialog Corporation at <<http://www.dialog.com>>.

relevant. Of these, 277 met the inclusion criteria. Team members also reviewed reference lists and consulted with other specialists in the field to identify relevant studies. All studies of economics, ethics, or feasibility that were applicable to the interventions under study were also examined. Among all the studies reviewed, 102 were excluded on the basis of limitations in their execution or design or because they duplicated information provided in an already included study. Excluded studies were not considered further. The remaining 175 studies were considered qualifying studies. The 12 Task Force recommendations in this report are based on the systematic review and evaluation of these qualifying studies, all of which had good or fair quality of execution.

On the basis of the evidence of effectiveness, the Task Force either strongly recommended or recommended 11 of the 12 interventions evaluated (Table 2). These 11 include four interventions to increase use of child safety seats (i.e., laws requiring use, communitywide information and enhanced enforcement campaigns, distribution and education programs, and incentive and education programs), three interventions to increase safety belt use (i.e., laws requiring use, primary enforcement laws, and enhanced enforcement programs), and four interventions to reduce alcohol-impaired driving (i.e., 0.08% blood alcohol concentration [BAC] laws, lower BAC limit laws for young [in the United States, those under the minimum legal drinking age] and inexperienced drivers, laws requiring a minimum drinking age of 21 years, and sobriety checkpoint programs). The Task Force found insufficient evidence on which to make a recommendation regarding the 12th intervention — education-only programs to improve child safety seat use — because of inconsistencies in the curricula, target populations, and effects of reported interventions.

In addition to these 12 interventions, reviews for two additional interventions to prevent motor-vehicle occupant injury — incentive programs to increase safety belt use and intervention training for servers of alcoholic beverages — are underway and will be included in a subsequent report.

USE OF THE RECOMMENDATIONS IN STATES AND COMMUNITIES

Given that motor-vehicle occupant injuries are the leading cause of death among persons aged 1–34 years in the United States (24), reducing the number of motor-vehicle crashes and crash-related occupant injuries should be relevant to most communities. States and communities can compare their current interventions and activities to prevent motor-vehicle injury with recommendations in this report, as well as with other relevant recommendations proposed by NHTSA (25), the National Transportation Safety Board* (NTSB) (26), DHHS (19), the American Medical Association (27), and the American Academy of Pediatrics (28,29).

The Task Force recommendations can be used to support or expand child safety seat distribution programs, bolster the use of incentives, and employ enhanced enforcement campaigns, all in conjunction with communitywide education efforts. For example, the recommendation for child safety seat distribution and education programs could help a community decide to concentrate the distribution of low-cost or no-cost child safety seats in low-income neighborhoods or to seek local sponsorship to defray the costs of seats distributed to needy families. In selecting and implementing interventions, communities should strive to develop a comprehensive program to reduce

*Available on the Internet at <<http://www.nts.gov>>.

TABLE 2. Recommendations from the Task Force on Community Preventive Services regarding the use of selected interventions to increase use of child safety seats, increase use of safety belts, and reduce alcohol-impaired driving

Intervention (No. of qualifying studies)	Task Force recommendation for use	Intervention description	Key findings* Median (range), number of studies
Interventions to increase child safety seat use			
Child safety seat law (n=9)	Strongly recommended	Requires infants and young children traveling in motor vehicles to be restrained in federally approved child safety seats appropriate for the child's age and size. State laws vary regarding the children to whom the laws apply (e.g., according to age, height, weight, seating position, or a combination of factors).	Child safety seat use: +13% (+5% to +35%), 3 studies. Fatal injuries: -35% (-57.3% to -25%), 3 studies. All fatal and nonfatal injuries: -17.3% (-35.9% to -10.5%), 5 studies.
Communitywide information and enforcement campaigns (n=4)	Recommended	Use media support and child safety seat displays in public sites to promote use. Use special enforcement strategies (e.g., checkpoints, dedicated law enforcement officials, or alternative penalties) to enforce existing child safety seat laws.	Child safety seat use: +12.3% (+3.8% to +20.8%), 5 studies.
Distribution and education programs (n=10)	Strongly recommended	Provide approved child safety seats to parents through loans, low-cost rentals, or giveaways. Include educational components of varying intensity.	Child safety seat use: +22.6% (+4% to +62.3%), 11 studies (range of follow-up times: immediate to 2 years).
Incentive and education programs (n=4)	Recommended	Provide rewards to children and parents for purchasing and correctly using child safety seats. Include educational components of varying intensity.	Child safety seat use: +9.9% (+4.8% to +36%), 6 studies (within first 5 months of program operation).
Education-only programs (n=6)	Insufficient evidence†	Provide information and teach skills to parents, children, or professional groups regarding the use of child safety seats.	Insufficient evidence to assess effectiveness in changing correct use or other outcomes. Three studies of perinatal education programs for parents regarding correct use of child safety seats.

TABLE 2. (Continued) Recommendations from the Task Force on Community Preventive Services regarding the use of selected interventions to increase use of child safety seats, increase use of safety belts, and reduce alcohol-impaired driving

Intervention (No. of qualifying studies)	Task Force recommendation for use	Intervention description	Key findings* Median (range), number of studies
			<p>One study of a preschool education program for children regarding correct use of child safety seats.</p> <p>Two studies of professional education — one for nurses on implementation of patient education programs and one for law enforcement officers on citation rates.</p>
Interventions to increase safety belt use			
Safety belt law (n=34)	Strongly recommended	<p>Requires the use of safety belts by motor-vehicle occupants not covered by the state’s child safety seat laws. Existing safety belt laws vary in their requirements. Most are age-based and cover occupants (usually front seat occupants) aged ≥16 years. These laws leave major gaps in coverage for occupants, primarily those aged 4–16 years.</p> <p>Specific requirements (e.g., age, seating position, fines, exceptions) vary by state.</p> <p>Studies assessed laws in the United States.</p>	<p>Observed safety belt use: +32% (+19.6% to +36.3%), 9 studies.</p> <p>Self-reported safety belt use: +15.8% (+13% to +18.7%), 4 studies.</p> <p>Police-reported safety belt use: +20.4% and +26%, 2 studies.</p> <p>Nonfatal injuries: -3.5% (-14.5% to +10.6%), 6 studies.</p> <p>Fatal injuries: -8.4% (-9% to -5%), 7 studies.</p> <p>Fatal and nonfatal injuries: -8.3% (-19.7% to -2.6%), 9 studies.</p>
Primary enforcement law (n=13)	Strongly recommended (instead of secondary law)	<p>Primary enforcement law allows a police officer to stop a vehicle solely for an observed belt law violation (without having other reasons for stopping the vehicle).</p> <p>Secondary enforcement law allows a police officer to issue a belt law citation only if the vehicle has been stopped for another violation.</p>	<p>Primary laws compared with secondary laws Observed safety belt use: +14.1% (+12% to +22.6%), 5 studies.</p> <p>Self-reported safety belt use: +22% and +1%, 2 studies.</p> <p>Fatal injuries: -7.7% (-13.9% to -3.1%), 3 studies.</p>

TABLE 2. (Continued) Recommendations from the Task Force on Community Preventive Services regarding the use of selected interventions to increase use of child safety seats, increase use of safety belts, and reduce alcohol-impaired driving

Intervention (No. of qualifying studies)	Task Force recommendation for use	Intervention description	Key findings* Median (range), number of studies
Enhanced enforcement (n=16)	Strongly recommended	Increased, rather than routine, enforcement at specific locations and times to target violations of safety belt laws. Media campaigns that publicize the enforcement activity are an important component.	Observed safety belt use: +17% (+8.3% to +24.0%), 16 studies. Fatal and nonfatal injuries: -6.7% and -15.3%, 2 studies.
Interventions to reduce alcohol-impaired driving			
0.08% blood alcohol con- centration (BAC) law (n=9)	Strongly recommended	Lowers the BAC at which it is illegal to drive a motor vehicle from 0.10 g/dL to 0.08 g/dL (0.08%).	Alcohol-related fatal crash outcomes: -7% (-15% to -4%), 25 measures.
Minimum legal drinking age (MLDA) (n=33)	Strongly recommended (i.e., maintaining MLDA at age 21 years)	Specifies an age below which the purchase or consumption of alcoholic beverages are not permitted.	<p>Among the targeted age group</p> <p><u>Raising the MLDA</u> Fatal crashes: -17% (-30% to -7%), 9 studies. Injury crashes: -15% (-33% to -6%), 4 studies. Other crashes: -21% and -18%, 2 studies. Fatal crash outcomes: -12% (-17% to -8%), estimated from 9 regression-based studies.</p> <p><u>Lowering the MLDA</u> Fatal crashes: +8% (+2% to +38%), 3 studies. Injury crashes: +5% (-2% to +22%), 4 studies. Other crashes: +22% and +186%, 2 studies.</p> <p>Insufficient evidence regarding the impact of raising the MLDA on alcohol-related crashes among adolescent drivers not directly affected by the law change.</p>
Lower BAC for young and inexperienced drivers (n=6)	Recommended	Establishes a separate, lower illegal BAC (usually ≥0.02 g/dL) for drivers targeted by the law. Studies assessed changes in state laws in the United States and Australia. U.S. laws apply to all drivers under the MLDA. In other countries, laws apply to either newly licensed drivers or newly licensed drivers under a specified age.	Alcohol-related fatal crashes: -17% (-24% to -9%), 3 studies. Injury crashes: -17% and -4%, 2 studies. Other crashes: -11%, 1 study.

TABLE 2. (Continued) Recommendations from the Task Force on Community Preventive Services regarding the use of selected interventions to increase use of child safety seats, increase use of safety belts, and reduce alcohol-impaired driving

Intervention (No. of qualifying studies)	Task Force recommendation for use	Intervention description	Key findings* Median (range), number of studies
Sobriety checkpoints (n=23)	Strongly recommended	<p>At random breath testing (RBT) checkpoints, all drivers stopped are given breath tests for BAC. Such checkpoints are not conducted in the United States.</p> <p>At selective breath testing (SBT) checkpoints, police must have reason to suspect the driver has been drinking before using breath tests.</p> <p>Media campaigns that publicize the enforcement activity are an important intervention component.</p>	<p>Random Breath Testing (RBT) Fatal crashes: -22% (-36% to -13%), 6 studies. Injury crash outcomes: -18% (-21% to -12%), 11 studies. Other crashes: -26% and -15%, 2 studies.</p> <p>Selective Breath Testing (SBT) Fatal crashes: -26% and -20%, 2 studies. Injury crashes: -21% (-24% to -5%), 6 studies. Other crashes: -24% (-35% to -13%), 5 studies.</p> <p>No discernible differences in effectiveness between RBT and SBT checkpoints.</p> <p>Similar magnitude of effects at long-term (>1 year) and short-term (≤1 year) follow-up.</p>

* Median effect sizes are a) absolute percentage point differences for all measures of child safety seat use and safety belt use and b) relative percent changes for all measures of injuries, crashes, and crash outcomes. When <7 studies are available, the range represents the entire range of effect measures identified; for ≥7 studies, an interquartile range is presented.

† A determination that evidence is insufficient should not be seen as evidence of ineffectiveness. A determination of insufficient evidence helps identify a) areas of uncertainty regarding effectiveness of an intervention and b) specific continuing research needs. In contrast, evidence of ineffectiveness leads to a recommendation that the intervention not be used.

motor-vehicle occupant injuries that includes legislation, enforcement, public education, training, and other community-oriented strategies. Improvements in each category will contribute to reductions in occupant-injury-related morbidity and mortality, and success in one area could contribute to improvements in the other areas.

The Task Force recommended or strongly recommended six state public health laws. Of these, three are in effect in all 50 states (i.e., laws requiring use of child safety seats, lower legal BAC for young or inexperienced drivers, and a minimum legal drinking age of 21 years). In addition, 49 states have laws requiring use of safety belts (New Hampshire has no such law).

Other laws reviewed by the Task Force were 0.08% BAC and primary enforcement safety belt laws. As of March 2001, 0.08% BAC laws had been enacted in 21 states, Washington, D.C., and Puerto Rico, and primary enforcement laws were in effect in 17 states, Washington, D.C., and Puerto Rico. In support of 0.08% BAC laws, the U.S. Congress included a provision in the 2001 Department of Transportation and Related Agencies Appropriations Act (30) requiring states to implement 0.08% BAC laws by fiscal year 2004 or risk losing federal highway construction funds.

The Task Force recommendations can be used to promote the adoption, maintenance, or strengthening of state or national laws or regulations. For example, at the state level, injury control program directors can use these recommendations to develop testimony regarding the evidence of effectiveness of different traffic safety laws for presentation to state legislatures. State legislators and their staff members can use the recommendations as they draft, debate, and vote on new or amended legislation. Advocacy and community groups can use the information to develop position statements regarding pending legislation at the state level. Health agencies can help educate the community regarding the importance and effectiveness of the laws and their enforcement.

Choosing effective interventions that are well-matched to state and local needs and capabilities, then carefully implementing those interventions, are vital steps in improving use of child safety seats and use of safety belts and in deterring alcohol-impaired driving. In setting priorities for the selection of interventions to meet local objectives, recommendations and other evidence provided in the *Community Guide* should be considered along with such local information as resource availability, administrative structures, and the economic, social, and regulatory environments of available organizations and practitioners. Involving other partners in these efforts could be useful. Examples of such partners are each state's Governor's Office of Highway Safety or local chapters of the National SAFE KIDS Campaign, available on the Internet at <<http://www.safekids.org>>; the National Safety Council, <<http://www.nsc.org>>; and Mothers Against Drunk Driving, <<http://www.madd.org>>. Additional information regarding applicability and economic information will be provided in the full report. Taking into consideration local goals and resources, the use of strongly recommended and recommended interventions should be given priority for implementation or enforcement.

ADDITIONAL INFORMATION REGARDING THE *COMMUNITY GUIDE*

During 2001–2002, *Community Guide* topics will be prepared and released as each is completed. Upcoming topics include diabetes, oral health, physical activity, sexual

behavior, cancer, and the sociocultural environment. A compilation of the recommendations and supporting evidence for these topics will be published in book form. Additional information regarding the Task Force and the *Community Guide* is available on the Internet at <<http://www.thecommunityguide.org>>.

References

1. CDC. Working to prevent and control injury in the United States: fact book for the year 2000. Atlanta, GA: US Department of Health and Human Services, CDC, 2000.
2. Rosenberg HM, Ventura SJ, Maurer JD, Heuser RL, Freedman MA. Births and deaths: United States, 1995. *Mon Vital Stat Rep* 1996;45(3, suppl 2):1–35.
3. Fingerhut LA, Warner M. Injury chartbook. Health, United States, 1996–97. Hyattsville, MD: US Department of Health and Human Services, CDC, 1997.
4. Sleet DA, Rosenberg ML. Injury control. In: Scutchfield FD, Keck CW, eds. Principles of public health practice. Albany, NY: Delmar Publishers, 1997:337–49.
5. National Highway Traffic Safety Administration. Traffic safety facts 1998: a compilation of motor vehicle crash data from the Fatality Analysis Reporting System and the General Estimates System. Washington, DC: US Department of Transportation, National Center for Statistics and Analysis, National Highway Traffic Safety Administration, 1999; Publication no. DOT HS 808-983.
6. Stussman BJ. National hospital ambulatory medical care survey: 1993 emergency department summary. *Adv Data* 1996;271:1–16. Available on the Internet at <<http://www.cdc.gov/nchs/data/ad/ad271.pdf>>. Accessed April 24, 2001.
7. Blincoe LJ. The economic cost of motor vehicle crashes, 1994. Washington, DC: National Highway Traffic Safety Administration, 1996; Publication no. DOT HS 808 425.
8. National Safety Council. Accident facts,™ 1998 ed. Itasca, IL: National Safety Council, 1998.
9. CDC. Motor-vehicle safety: a 20th century public health achievement. *MMWR* 1999;48:369–74.
10. CDC. Prevention of motor vehicle-related injuries: a compendium of articles from the *Morbidity and Mortality Weekly Report* 1985–1996. Atlanta, GA: US Department of Health and Human Services, CDC, 1997.
11. Sleet DA. Reducing motor vehicle trauma through health promotion programming. *Health Educ Q* 1984;11:113–25.
12. Task Force on Community Preventive Services. Recommendations regarding interventions to improve vaccination coverage in children, adolescents, and adults. *Am J Prev Med* 2000;18(1S):92–6.
13. CDC. Vaccine-preventable diseases: improving vaccination coverage in children, adolescents, and adults—a report on recommendations of the Task Force on Community Preventive Services. *MMWR* 1999;48(No. RR-8).
14. Briss PA, Rodewald LE, Hinman AR, et al. Reviews of evidence regarding interventions to improve vaccination coverage in children, adolescents, and adults. *Am J Prev Med* 2000;18(1S):97–140.
15. Task Force on Community Preventive Services. Recommendations regarding interventions to reduce tobacco use and exposure to environmental tobacco smoke. *Am J Prev Med* 2001;20(2S):10–5.
16. CDC. Strategies for reducing exposure to environmental tobacco smoke, increasing tobacco-use cessation, and reducing initiation in communities and health-care systems: a report on recommendations of the Task Force on Community Preventive Services. *MMWR* 2000;49(No. RR-12).
17. Hopkins DP, Briss PA, Ricard CJ, et al. Reviews of evidence regarding interventions to reduce tobacco use and exposure to environmental tobacco smoke. *Am J Prev Med* 2001;20(2S):16–66.
18. Briss PA, Zaza S, Pappaioanou M, et al. Developing an evidence-based *Guide to Community Preventive Services*—methods. *Am J Prev Med* 2000;18(1S):35–43.
19. US Department of Health and Human Services. Healthy People 2010, 2nd ed. With understanding and improving health and objectives for improving health. 2 vols. Washington, DC: US Government Printing Office, 2000.

20. National Highway Traffic Safety Administration. Traffic safety facts 1999: children. Washington, DC: US Department of Transportation, National Highway Traffic Safety Administration, 2000; publication no. DOT HS 809 087.
21. National Highway Traffic Safety Administration. Traffic safety facts 1999: occupant protection. Washington, DC: US Department of Transportation, National Highway Traffic Safety Administration, 2000; publication no. DOT HS 809 090.
22. Zador PL, Krawchuk SA, Voas RB. Alcohol-related relative risk of driver fatalities and driver involvement in fatal crashes in relation to driver age and gender: an update using 1996 data. *J Stud Alcohol* 2000;61:387-95.
23. Zaza S, Wright-De Agüero LK, Briss PA, et al. Data collection instrument and procedure for systematic reviews in the *Guide to Community Preventive Services*. *Am J Prev Med* 2000;18(1S):44-74.
24. Pickle LW, Mungiole M, Jones GK, White AA. Atlas of United States mortality. Hyattsville, MD: US Department of Health and Human Services, Public Health Service, CDC, 1996; DHHS publication no. (PHS) 97-1015.
25. National Highway Traffic Safety Administration. Highway safety program advisories. Washington, DC: US Department of Transportation, National Highway Traffic Safety Administration, 1990; Publication no. DOT HS 807 655.
26. National Transportation Safety Board. We are all safer: NTSB-inspired improvements in transportation safety. 2nd ed. Washington, DC: National Transportation Safety Board. Available on the Internet at <<http://www.nts.gov/Publictn/1998/SR9801.pdf>>. Accessed January 25, 2001.
27. American Medical Association. Operating vehicles under the influence of alcohol or other drugs/underage drinking and driving. Available on the Internet at <<http://www.ama-assn.org/special/aos/alcohol1/policy/drinking.htm>>. Accessed December 26, 2000.
28. American Academy of Pediatrics. Selecting and using the most appropriate car safety seats for growing children: guidelines for counseling parents (RE9618). Available on the Internet at <<http://www.aap.org/policy/01352.html>>. Accessed December 26, 2000.
29. American Academy of Pediatrics. The teenage driver (RE9642). Available on the Internet at <<http://www.aap.org/policy/re9643.html>>. Accessed December 26, 2000.
30. Department of Transportation and Related Agencies Appropriations Act, 2001. Pub L No. 106-346, 114 Stat 1356.

Use of trade names and commercial sources is for identification only and does not imply endorsement by the U.S. Department of Health and Human Services.

References to non-CDC sites on the Internet are provided as a service to *MMWR* readers and do not constitute or imply endorsement of these organizations or their programs by CDC or the U.S. Department of Health and Human Services. CDC is not responsible for the content of pages found at these sites.

MMWR

The *Morbidity and Mortality Weekly Report (MMWR)* Series is prepared by the Centers for Disease Control and Prevention (CDC) and is available free of charge in electronic format and on a paid subscription basis for paper copy. To receive an electronic copy on Friday of each week, send an e-mail message to listserv@listserv.cdc.gov. The body content should read *SUBscribe mmwr-toc*. Electronic copy also is available from CDC's World-Wide Web server at <http://www.cdc.gov/mmwr/> or from CDC's file transfer protocol server at <ftp://ftp.cdc.gov/pub/Publications/mmwr/>. To subscribe for paper copy, contact Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402; telephone (202) 512-1800.

Data in the weekly *MMWR* are provisional, based on weekly reports to CDC by state health departments. The reporting week concludes at close of business on Friday; compiled data on a national basis are officially released to the public on the following Friday. Address inquiries about the *MMWR* Series, including material to be considered for publication, to: Editor, *MMWR* Series, Mailstop C-08, CDC, 1600 Clifton Rd., N.E., Atlanta, GA 30333; telephone (888) 232-3228.

All material in the *MMWR* Series is in the public domain and may be used and reprinted without permission; citation as to source, however, is appreciated.